Aetiological investigations for children and young people with permanent hearing loss : are the current guidelines being followed?

Dr Vishakha Phatak, Consultant Paediatrician Bradford Teaching Hospitals Foundation Trust



Jan 2016 BAPA Annual Conference

Aim:

Evaluate local practice for offering Level 1 investigations

Gold standard:

BAPA and BAAP guidelines (2008-2009)



BAPA and BAAP guidelines:

	CMV	Vision	Urine dipstick (Alport)	Family	Connexin	ECG
Bilateral						
Mild	\checkmark	\checkmark	\checkmark	\checkmark	-	
Mod	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	
Severe- profound	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
Unilateral	\checkmark	\checkmark		\checkmark		



Methodology:

Period: March 2013- February 2014

All new referrals

Data collection: October 2014 to January 2015




Methodology :

Information collected:

- Age
- Severity of hearing loss
- Unilateral/bilateral
- Investigations offered-accepted and rejected



Sample:




Distribution of hearing loss:

	Mild	Moderate	Severe- profound	Total
Bilateral	10	14	7	31
Unilateral	4	3	3	11
	14 (33%)	17 (40%)	11(26%)	42




Results:




Investigations offered (% of total)



Investigations







Majority but not all offered Level 1 investigations

Area for further work identified

- documentation of reasons for refusal



Recommendations:

Re-audit after revised guidelines published

Multi-centric study

Research reasons for refusal of investigations



Thank you

